

DOI: 10.11931/guihaia.gxzw202312059

温放, 韦毅刚, 李晓娟, 等, 2024. 辐冠苣苔属(苦苣苔科)一新替代名称——*Radiaticorollarus* Y. G. Wei, F. Wen & Lei Cai [J]. 广西植物, 44(4): 619–620.

WEN F, WEI YG, LI XJ, et al., 2024. *Radiaticorollarus*, a new replacement name for *Actinostephanus* (Gesneriaceae, Didymocarpoideae) [J]. Guihaia, 44(4): 619–620.



***Radiaticorollarus*, a new replacement name for *Actinostephanus* (Gesneriaceae, Didymocarpoideae)**

WEN Fang^{1,2}, WEI Yigang^{1,2}, LI Xiaojuan^{1,2}, XIONG Chi^{1,2}, CAI Lei^{3*}

(1. Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin 541006, Guangxi, China; 2. National Gesneriaceae Germplasm Resources Bank of GXIB, Gesneriad Committee of China Wild Plant Conservation Association, Gesneriad Conservation Center of China, Guilin Botanical Garden, Chinese Academy of Sciences, Guilin 541006, Guangxi, China; 3. Yunnan Key Laboratory for Integrative Conservation of Plant Species with Extremely Small Populations, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China)

Abstract: *Actinostephanus* F. Wen, Y. G. Wei & L. F. Fu is a monotypic genus in Gesneriaceae endemic to Guangdong, China. It is named for its radiate corolla lobes, the relatively rare character among Gesneriaceae. Some scholars have suggested that the name *Actinostephanus* is easily confused with the generic name of the extinct fossil phytoplankton *Actinostephanos* Khursevich, and proposed that *Actinostephanus* should be treated as a later homonym of *Actinostephanos*. According to the *International Code of Nomenclature of Algae, Fungi and Plants (Shenzhen Code)* (2018), we here propose a new name *Radiaticorollarus* Y. G. Wei, F. Wen & Lei Cai for the later homonym *Actinostephanus* F. Wen, Y. G. Wei & L. F. Fu, and treat *Actinostephanus enpingensis* F. Wen, Y. G. Wei & Z. B. Xin as a synonym of the new combination *Radiaticorollarus enpingensis* (F. Wen, Y. G. Wei & Z. B. Xin) F. Wen, Y. G. Wei & Lei Cai.

Key words: *Actinostephanus*, *Actinostephanus enpingensis*, Gesneriaceae, new name, new combination

CLC number: Q941 **Document code:** A **Article ID:** 1000-3142(2024)04-0619-02

辐冠苣苔属(苦苣苔科)一新替代名称 ——*Radiaticorollarus* Y. G. Wei, F. Wen & Lei Cai

温 放^{1,2}, 韦毅刚^{1,2}, 李晓娟^{1,2}, 熊 驰^{1,2}, 蔡 磊^{3*}

(1. 广西喀斯特植物保育与恢复生态学重点实验室, 广西壮族自治区广西植物研究所, 广西桂林 541006; 2. 广西植物研究所国家苦苣苔科种质资源库, 中国野生植物保护协会苦苣苔专委会, 中国苦苣苔科植物保育中心, 中国科学院桂林植物园, 广西桂林 541006; 3. 中国科学院昆明植物研究所, 云南省极小种群野生植物综合保护重点实验室, 昆明 650201)

收稿日期: 2023-12-31 接受日期: 2024-01-24

基金项目: 广西科技计划项目(桂科ZY21195050); 中国科学院战略生物资源能力建设项目(KFJ-BRP-017-68)。

第一作者: 温放(1976—), 博士, 研究员, 主要从事园林植物和观赏植物育种和应用、植物分类学和保育生物学研究,(E-mail) wenfang760608@139.com。

*通信作者: 蔡磊, 博士, 副研究员, 主要从事极小种群野生植物的保护生物学、植物分类学和植物资源学研究,(E-mail) cailei@mail.kib.ac.cn。

摘要:辐冠苣苔属为中国广东特有苦苣苔科单型属,因其具有苦苣苔科中相对比较少见的辐射状花冠裂片而得名。有学者提出因其属名 *Actinostephanus* F. Wen, Y. G. Wei & L. F. Fu 和已灭绝的化石浮游植物属名 *Actinostephanos* Khursevich 极其相似而易于混淆,应处理为晚出同名。依据 2018 年《国际藻类、菌物和植物命名法规(深圳法规)》,该文为该晚出同名拟定了一个新名称,即 *Radiaticorollarus* Y. G. Wei, F. Wen & Lei Cai, 并将其属下辐冠苣苔 *Actinostephanus enpingensis* F. Wen, Y. G. Wei & Z. B. Xin 处理为新组合 *Radiaticorollarus enpingensis* (F. Wen, Y. G. Wei & Z. B. Xin) F. Wen, Y. G. Wei & Lei Cai 的异名。

关键词:辐冠苣苔属,辐冠苣苔,苦苣苔科,新名称,新组合

The genus *Actinostephanus* F. Wen, Y. G. Wei & L. F. Fu was named for its radiate corolla lobes, the relatively rare character among Gesneriaceae. It includes a single species, *A. enpingensis* F. Wen, Y. G. Wei & Z. B. Xin. The genus name, “*Actinostephanus*”, is derived from Greek, with the prefix “*Actino-*” coming from ἀκτίς (Aktís, meaning “ray” or “radiate”) and “*stephanos*” from Στέφανος (Stéphanos, meaning “crown”). This type of corolla characteristic is rare in Chinese Gesneriaceae. Before the discovery of this new genus, only three species, belonging to two different genera, have radially spreading corollas: *Bournea sinensis* Oliv., *B. leiophylla* (W. T. Wang) W. T. Wang & K. Y. Pan ex W. T. Wang, and *Oreocharis esquirolii* H. Lév.

However, Khursevich (1989) had named a kind of extinct fossil phytoplankton *Actinostephanos* Khursevich (1989: 487) using a homological name. Despite the difference of a single letter between the two names, “u” and “o”, which is sufficient to differentiate them, Deshmukh (2022: 1127) believed they are extremely similar and prone to confusion and should be treated as homonyms, thus requesting a binding decision that the names *Actinostephanus* Wen et al. and *Actinostephanos* Khursevich are easily confused and should be treated as homonyms. This request was recommended by The Nomenclature Committee for Algae (Anderson, 2023) and approved by General Committee.

Therefore, according to Article 53.1 of the *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code)* (Turland et al., 2018), a new name, *Radiaticorollarus* Y. G. Wei, F. Wen & Lei Cai, is proposed herein for the later homonym *Actinostephanus* F. Wen, Y. G. Wei & L. F. Fu. *Radiaticorollarus* is composed of two Latin words, “*Radiati-*” deriving from “*Radiatus*”, meaning radiate, and “*corollarus*” from “*corolla*”, meaning corolla. This replacement name retains the same meaning as the original *Actinostephanus*, referring to the unique species in this genus known to have a campanulate corolla tube and radially symmetrical corolla lobes, distinct from all other known genera in China.

Taxonomic treatment

Radiaticorollarus Y. G. Wei, F. Wen & Lei Cai, nom. nov.

Replaced Synonym: — *Actinostephanus* F. Wen,

Y. G. Wei & L. F. Fu, *PhytoKeys* 193: 95, 2022, nom. illeg., non *Actinostephanos* Khursevich, *Bot. Zhurn.* 74: 487 1989 (Algae, Bacillariophyta). **TYPE:** — *Radiaticorollarus enpingensis* (F. Wen, Y. G. Wei & Z. B. Xin) F. Wen, Y. G. Wei & Lei Cai (≡ *Actinostephanus enpingensis* F. Wen, Y. G. Wei & Z. B. Xin).

Radiaticorollarus enpingensis (F. Wen, Y. G. Wei & Z. B. Xin) F. Wen, Y. G. Wei & Lei Cai, comb. nov.

Basionym: — *Actinostephanus enpingensis* F. Wen, Y. G. Wei & Z. B. Xin, *PhytoKeys* 193: 96. 2022. **TYPE:** — CHINA. Guangdong Province: Enping City, Naji Town, Qixingkeng provincial natural reserve, growing on the cliffs or slopes of montane yellow soil and sandy loam near a stream, 22°11.1808' N, 112°5.841' E, ca. 153 m, 19 May 2021, Chen Xiaoyun & Liang Junjie 210519 - 01 (holotype: IBK, isotypes: IBK).

Acknowledgments

We would like to thank Research Professor DENG Yunfei of the South China Botanical Garden, Chinese Academy of Sciences for his suggestions and strong support. We also thank the editors and anonymous reviewers for their positive comments and help on the manuscript.

References:

- ANDERSEN RA, 2023. Report of the Nomenclature Committee for Algae: 25. [J]. *Taxon*, 72(6): 1355.
- DESHMUKH UB, 2022. (104) Request for a binding decision on whether the names *Actinostephanos* Khursevich (Bacillariophyta) and *Actinostephanus* F. Wen & al. (Gesneriaceae) are sufficiently alike to be confused [J]. *Taxon*, 71(5): 1127.
- KHURSEVICH GK, 1989. New data on morphology and systematic position of *Stephanodiscus matrensis* (Bacillariophyta) [J]. *Bot Zh*, 74(4): 487–489.
- TURLAND NJ, WIERSEMA JH, BARRIE FR, et al., 2018. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017 [M]. Koeltz botanical books, Glashütten. *Regnum Vegetabile* 159. <https://doi.org/10.12705/Code.2018>.
- WEN F, XIN ZB, HONG X, et al., 2022. *Actinostephanus* (Gesneriaceae), a new genus and species from Guangdong, South China [J]. *PhytoKeys*, 193: 89–106.

(责任编辑 蒋巧媛 王登惠)